


Dual prescription refill information



Patient name _____

Prescription number _____

Owner's name _____

Treatment start date _____

Veterinarian _____

Dual prescription refill schedule

| DAY | MIX | DATE ADMINISTERED | SCHEDULED DOSE | ACTUAL DOSE | REACTIONS | CURRENT ITCHINESS LEVEL (Scale: 10 = high, 1 = none) |
|-----|-------|-------------------|--|-------------|-----------|--|
| 1 | Mix 1 | | $\frac{1}{4}$ of previous tolerated dose | | | |
| 3 | Mix 2 | | $\frac{1}{4}$ of previous tolerated dose | | | |
| 8 | Mix 1 | | $\frac{1}{2}$ of previous tolerated dose | | | |
| 10 | Mix 2 | | $\frac{1}{2}$ of previous tolerated dose | | | |
| 15 | Mix 1 | | $\frac{3}{4}$ of previous tolerated dose | | | |
| 17 | Mix 2 | | $\frac{3}{4}$ of previous tolerated dose | | | |
| 29 | Mix 1 | | Previous tolerated dose | | | |
| 31 | Mix 2 | | Previous tolerated dose | | | |
| 50 | Mix 1 | | Previous tolerated dose | | | |
| 52 | Mix 2 | | Previous tolerated dose | | | |

READ THE FOLLOWING INFORMATION BEFORE USING THIS VIAL

Allergenic extracts tend to lose potency over time. A newly prepared prescription is likely stronger than the one formulated for your pet a few months ago. To compensate for the change in strength of the new prescription vial, the pet should receive a lower volume for the first month to avoid any reaction. If your pet has a reaction at any time during this build-up period, do not continue to increase the volume. Call your veterinarian to help you adjust the schedule. Start the new schedule at the start of your next routinely scheduled dose.

Return to the volume and frequency schedule previously found as best suited to the pet's response. Order refills from your veterinarian a minimum of 6 weeks in advance. Always have this schedule updated and in hand when contacting or visiting your veterinarian. Please consult your veterinarian before making any dosage changes.

**Reviewed and approved
by a veterinary dermatologist.**

IDEXX

STALLERGENES  GREER

Please see reverse side for more information.

Dual prescription refill dosage schedule

Understanding the dosage schedule

The dosage schedule tells you which vials to use, how much to inject, and when to give the injections. The Build-up Phase increases the dose gradually. After increasing to the maximum tolerated dose, the pet is “maintained” at one dose given every 10–21 days. This is the Maintenance Dose.

Since you are receiving a refill prescription, you have already been through the build-up phase and have reached the maintenance phase. It is important to get your pet back to their maintenance dose (previously tolerated dose) and the injection schedule you have found to be optimal. However, because allergenic extracts tend to lose potency over time, it is important to revisit the build-up phase. Please use the guide on the opposite page to help you, but always consult your veterinarian if you have any concerns.

Example:

| DAY | MIX | PREVIOUSLY TOLERATED DOSE | BUILD-UP ADJUSTMENT | DOSE |
|-----|-----|---------------------------|---------------------|---------|
| 1 | 1 | 1.0 mL | 0.25 x 1.0 mL | 0.25 mL |
| 3 | 2 | 1.0 mL | 0.25 x 1.0 mL | 0.25 mL |
| 8 | 1 | 1.0 mL | 0.50 x 1.0 mL | 0.50 mL |
| 10 | 2 | 1.0 mL | 0.50 x 1.0 mL | 0.50 mL |
| 15 | 1 | 1.0 mL | 0.75 x 1.0 mL | 0.75 mL |
| 17 | 2 | 1.0 mL | 0.75 x 1.0 mL | 0.75 mL |
| 29 | 1 | 1.0 mL | 1.0 x 1.0 mL | 1.0 mL |

Success depends on finding what dosage and frequency of doses works best for your pet. Most pets do not follow the dosage schedule exactly and this is offered as a guide.

Refer to the **GREER® Aller-g-complete® Guide** for more information.

Please consult your veterinarian before making any dosage changes.