


Find what the microscope misses.¹⁻⁵

Don't rely on eggs alone. Fecal Dx™ antigen testing detects antigens secreted from the infecting worm.

- + Identify infections prior to eggs being shed.
- + Avoid false-negative results caused by intermittent shedding of eggs.
- + Detect worms even if eggs are not present in the fecal specimen.

Help prevent spread of disease to family members and pets.

Detecting infections sooner helps you treat effectively, reducing the risk of infection to other pets and family members.⁶



A single test finds hookworms, roundworms, whipworms, and flea tapeworms.



Earlier detection of hookworms, roundworms, and whipworms than fecal flotation.¹⁻⁵




Detects 5X more infections than fecal flotation alone.^{3,5,7}


Set your sights on parasites.

Fecal Dx[®] antigen testing helps you detect more parasites earlier than ever before.

Detect more. Results indicate that adding Fecal Dx antigen testing to your fecal flotation testing may double the number of parasite infections detected.⁸



Detect earlier. Because of the lack of egg detection with fecal flotation testing during the prepatent period and single-sex infections, many parasite infections may go undetected for a period of time.¹



*Data represents fecal testing performed during wellness visits.

Parasites are everywhere.


U.S. dog park study confirms the need to test and treat even well-cared for dogs.⁹

- + Approximately 3,000 fresh canine fecal specimens were collected.
- + 288 dog parks in 30 metropolitan areas in the U.S. were included.
- + Fecal Dx™ antigen testing, *Giardia* antigen testing, and fecal flotation by centrifugation were conducted.

Key results:



85% of parks tested positive for an intestinal parasite.



1 in 5 dogs tested positive for intestinal parasites.

In a similar study performed in Alberta, Canada, involving 774 participants with fecal specimens being collected from 7 dedicated off-leash dog parks, the overall prevalence of canine intestinal parasites in client-owned dogs was similar to numbers of parasite infections from dog parks in the northwestern U.S.¹⁰



50% of dog parks tested positive for hookworms, roundworms, or whipworms.



Test to protect with Fecal Dx antigen testing:



Fecal Dx™ antigen testing catches infections missed by the microscope.¹⁻⁴




Parasite infections may go undetected, even in dogs receiving preventives.⁹




Ideally, dogs are protected by broad-spectrum, year-round prevention.

Only IDEXX Reference Laboratories offers a complete fecal testing solution:



Zinc sulfate centrifugation.



Egg and parasite identification by expert technicians.



Fecal Dx antigen testing.

Common test codes for fecal testing.

For healthy dogs and cats:

Fecal Dx™ Antigen Panel (test code FDX)

Hookworm, roundworm, whipworm, and flea tapeworm antigens by ELISA

For sick dogs and cats:

Fecal Dx™ Antigen Profile, with *Giardia* (test code FDXGP)

Fecal flotation, *Giardia*, hookworm, roundworm, whipworm, and flea tapeworm antigens by ELISA

Expanded options for puppies and kittens:

Fecal Dx™ Profile (test code FDXP)

Fecal flotation, hookworm, roundworm, whipworm, and flea tapeworm antigens by ELISA.

Also available as home collection kit:

IDEXX FecalChek®

Giardia, hookworm, roundworm, whipworm, and flea tapeworm antigens by ELISA.

Available to order via IDEXX Online Orders.

**Improved turnaround time:
1 day for most customers.***

*1-2 days for customers based in Alberta.

Scoop. Seal. Send. It's that simple.

The IDEXX FecalChek Home Collection Kit makes collection for pet owners a snap.

To make it easier for owners—and to make sure more pets get screened for intestinal parasites—we created IDEXX FecalChek® Home Collection Kit. It has everything pet owners need to send a specimen to the reference lab themselves. They simply scoop up the specimen at home, seal it in the vial provided, and then send it directly to our reference lab in a preaddressed envelope. Your practice will receive test results as expected with any IDEXX fecal test offering.

The home collection kit can increase compliance by up to 20%,² which means:



Healthy pets

With early detection, Fecal Dx™ antigen testing can lead to earlier treatment, which may reduce the risk of infection to other pets and family members, too.⁶



Happy clients

It's easy and convenient for pet owners. They simply collect the specimen and drop the kit into nearest mailbox.



Profitable practice

Home collection enhances your clients' experience while creating opportunity for increased diagnostic and treatment revenue.

Order your kits at order.idexx.ca




Just start with VetConnect PLUS, your go-to diagnostic resource for efficiency and flexibility.

VetConnect® PLUS offers the diagnostic tools you need for fast, effective clinical decision-making.




Spot trends and abnormalities with at-a-glance patient historical results.

Easily compare results side-by-side or graph patient historical results for visual trend analyses.




Improve client communications and increase compliance.

Share diagnostic results and personalized client-friendly result summaries with one click.



Access differential diagnoses to help interpret results and determine next steps

Click to view guidance from IDEXX board-certified specialists using the latest industry protocols, research, and publications.



Get results in 1–2 days for Fecal Dx™ antigen testing.

Results are automatically added to the patient history in VetConnect PLUS.

To activate your VetConnect PLUS account, please complete the online registration form on idexx.ca/vetconnectplus. Sign in at vetconnectplus.ca.

Manage patient diagnostics from your smartphone, tablet, browser, or practice management system.

Mobile app is available for iPhone® and Android™ devices.



References

1. Elsemore DA, Geng J, Flynn L, Cruthers L, Lucio-Forster A, Bowman DD. Enzyme-linked immunosorbent assay for coproantigen detection of *Trichuris vulpis* in dogs. *J Vet Diagn Invest.* 2014;26(3):404–411.
2. Data on file at IDEXX Laboratories, Inc. Westbrook, Maine USA.
3. Data on file at IDEXX Laboratories, Inc. Westbrook, Maine USA. Aggregate detection of hookworm, roundworm, and whipworm infections.
4. Elsemore DA, Geng J, Cote J, Hanna R, Lucio-Forster A, Bowman DD. Enzyme-linked immunosorbent assays for coproantigen detection of *Ancylostoma caninum* and *Toxocara canis* in dogs and *Toxocara cati* in cats. *J Vet Diagn Invest.* 2017;29(5):645–653. doi:10.1177/1040638717706098
5. Elsemore D, Beall M, Bezold T, et al. Detection of *Dipylidium caninum* coproantigen in experimental and natural infections [AAVP Abstract 23]. Paper presented at: American Association of Veterinary Parasitologists 67th Annual Meeting; June 26, 2022; Snowbird, UT
6. CAPC guidelines: controlling internal and external parasites in U.S. dogs and cats. Companion Animal Parasite Council. Accessed March 9, 2023. www.petsandparasites.org/resources/capc-guidelines
7. Data on file at IDEXX Laboratories, Inc. Westbrook, Maine USA. Preliminary data on positive rate for *Dipylidium caninum* is approximately 6%.
8. Sweet S, Hegarty E, McCrann DJ, Coyne M, Kincaid D, Szlosek D. A 3-year retrospective analysis of canine intestinal parasites: fecal testing positivity by age, U.S. geographical region and reason for veterinary visit. *Parasit Vectors.* 2021;14(1):173. doi:10.1186/s13071-021-04678-6
9. Stafford K, Kollasch TM, Duncan KT, et al. Detection of gastrointestinal parasitism at recreational canine sites in the USA: the DOGPARGS study. *Parasit Vectors.* 2020;13(1):275. doi:10.1186/s13071-020-04147-6
10. Visscher D, Porter E, Sweet S, Szlosek D, Horr S. Canine nematode and *Giardia* spp. infections in dogs in Edmonton, Alberta, the “CANIDA” study. *Parasit Vectors.* 2022;15(1):294. doi:10.1186/s13071-022-05386-5